

D7



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) EP 1 022 332 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:
26.07.2000 Bulletin 2000/30(51) Int. Cl.⁷: C12N 9/10, C12N 15/54,
C12N 1/19, C12P 19/00

(21) Application number: 98941819.9

(86) International application number:
PCT/JP98/04087

(22) Date of filing: 10.09.1998

(87) International publication number:
WO 99/13059 (18.03.1999 Gazette 1999/11)(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: 10.09.1997 JP 24515497

(71) Applicant:
Meiji Seika Kaisha, Ltd.
Tokyo 104-8002 (JP)

- NAKANE, Akitaka,
Meiji Seika Kaisha, Ltd.
Sakado-shi, Saitama 350-0289 (JP)
- KONO, Toshiaki,
Meiji Seika Kaisha, Ltd.
Sakado-shi, Saitama 350-0289 (JP)


(72) Inventors:
• YANAI, Koji,
Meiji Seika Kaisha, Ltd.
Sakado-shi, Saitama 350-0289 (JP)(74) Representative:
Lewin, John Harvey et al
Elkington and Fife,
Prospect House,
8 Pembroke Road
Sevenoaks, Kent TN13 1XR (GB)(54) **BETA-FRUCTOFURANOSIDASE AND GENE THEREOF**(57) A novel β -fructofuranosidase and its gene are disclosed. A polypeptide comprising the amino acid sequence of SEQ ID No. 1 or No. 3 is an enzyme having β -fructofuranosidase activity and high transferase activity, and is capable of efficiently producing fructooligosaccharides.

FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D

EP 1 022 332 A1

Description

Background of the Invention5 Field of the Invention

[0001] The present invention relates to a β -fructofuranosidase having a fructose transferase activity, which is useful for the industrial production of fructooligosaccharides, and its gene and use.

10 Background Art

[0002] The molecular structure of a fructooligosaccharide is the same as that of sucrose, except that the fructose half of a fructooligosaccharide is coupled with another one to three fructose molecules at positions C1 and C2 via a β -bond. Fructooligosaccharides are indigestible sugars known for their physiological advantages, such as the facilitation of Bifidobacterial growth in the intestines, metabolic stimulation for cholesterol and other lipids, and little cariogenicity.

[0003] Fructooligosaccharides are found in plants, such as asparagus, onion, Jerusalem-artichoke and honey. They are also synthesized from sucrose by the newly industrialized mass production technique using fructosyltransferase reaction which is catalyzed by a β -fructofuranosidase derived from a microorganism.

[0004] The molecular structure of 1-kestose and nystose, which make up component of industrially produced fructooligosaccharide mixtures of today, are the same as that of sucrose except that their fructose half is coupled with one and two molecules of fructose, respectively. It has been found recently that their high-purity crystals exhibit new desirable characteristics both in physical properties and food processing purpose while maintaining the general physiological advantages of fructooligosaccharides (Japanese Patent Application No. 222923/1995, Japanese Laid-Open Publication No. 31160/1994). In this sense, they are fructooligosaccharide preparations having new features.

[0005] In consideration of the above, some of the inventors have already proposed an industrial process for producing crystal 1-kestose from sucrose (Japanese Patent Application No. 64682/1996, Japanese Patent Application No. 77534/1996, and Japanese Patent Application No. 77539/1996). According to this process, a β -fructofuranosidase harboring fructosyltransferase activity is first allowed to act on sucrose to produce 1-kestose; the resultant 1-kestose is fractionated to a purity of 80% or higher by chromatographic separation; then, using this fraction as a crystallizing sample, crystal 1-kestose is obtained at a purity of 95% or higher. The β -fructofuranosidase harboring fructosyltransferase activity used in this process should be able to produce 1-kestose from sucrose at a high yield while minimizing the byproduct nystose, which inhibits the reactions in the above steps of chromatographic separation and crystallization. In the enzyme derived from Aspergillus niger, which is currently used for the industrial production of fructooligosaccharides mixtures, the 1-kestose yield from sucrose is approximately 44%, while 7% is turned to nystose (Japanese Patent Application No. 64682/1996). These figures suggest that the enzyme has room for improvement in view of the industrial production of crystal 1-kestose.

[0006] As a next step, some of the inventors have successfully screened new enzymes having more favorable characteristics from Penicillium roqueforti and Scopulariopsis brevicaulis. These enzymes were able to turn 47% and 55% of sucrose into 1-kestose, respectively, and 7% and 4% to nystose (Japanese Patent Application No. 77534/1996, and Japanese Patent Application No. 77539/1996). These enzymes are inferior in productivity and stability to the enzyme derived from Aspergillus niger, and have room for improvement in view of the industrial production of crystal 1-kestose.

[0007] Thus, some of the inventors had paid attention to the procedure of genetic engineering as a process for improving the productivity of the enzyme, isolated the gene encoding β -fructofuranosidase from Penicillium roqueforti and Scopulariopsis brevicaulis, respectively, and conducted the structure analysis (PCT/JP97/00757). As a result, the translation regions encoding 565 amino acids and 574 amino acids as a mature protein were respectively deduced in the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis and their expression products were shown to have β -fructofuranosidase activity, like the β -fructofuranosidase gene from Aspergillus niger (L.M. Boddy et al., Curr. Genet., 24, 60-66 (1993)).

50 Summary of the Invention

[0008] The inventors have now found that the addition of 38 and 39 amino acids to the C-terminal of the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis, which were previously found by some of the inventors, improves its activity.

[0009] Thus, an object of the present invention is to provide a novel β -fructofuranosidase and its gene.

[0010] The novel β -fructofuranosidase according to the present invention is a polypeptide comprising the amino acid sequence of SEQ ID No. 1 or 3 or a homologue thereof.

[0011] Furthermore, the gene according to the present invention is a DNA encoding the above polypeptide.

EP 1 022 332 A1

[0012] The amino acid sequence of SEQ ID No. 1 or 3 according to the present invention is constructed by adding 38 and 39 amino acids to the C-terminals of the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis, which were previously found by some of the inventors as described above. It has been found that an intron actually exists at the region of the β -fructofuranosidase gene, which was presumed to encode the C-terminal amino acids by some of the present inventors and that the β -fructofuranosidase genes further encode 38 and 39 amino acids of the C-terminal. Surprisingly, the β -fructofuranosidase activity was remarkably improved by adding these amino acids to the C-terminal, as compared with the protein to which these sequences are not added.

Brief Description of the Drawing

[0013]

Figures 1A, B, C and D show the construction of expression vector pYPEN02 in which a gene encoding the enzyme protein consisting of the amino acid sequence of SEQ ID No. 1 is introduced, and expression vector pYPEN01 in which a gene encoding the enzyme protein consisting of the amino acid sequence from 1 to 565 of amino acid sequence of SEQ ID No. 1 is introduced.

Figures 2A and B show the construction of expression vector pYSCOP02 in which a gene encoding the enzyme protein consisting of the amino acid sequence of SEQ ID No. 3 is introduced, and expression vector pYSCOP01 in which a gene encoding the enzyme protein consisting of the amino acid sequence from 1 to 574 of amino acid sequence of SEQ ID No. 3 is introduced.

Detailed Description of the Invention β -fructofuranosidase

[0014] The polypeptide according to the present invention comprises the amino acid sequence of SEQ ID No. 1 or 3. This polypeptide having the amino acid sequence of SEQ ID No. 1 or 3 has enzymatic activity as β -fructofuranosidase. The polypeptide according to the present invention involves a homologue of the amino acid sequence of SEQ ID No. 1 or 3 as shown in the sequence listing. The term "homologue" refers to an amino acid sequence in which one or more amino acids (for example, one to several amino acids) are inserted, substituted or deleted in, or added to either or both of the terminals of, the amino acid sequence of SEQ ID Nos. 1 and 3 while retaining β -fructofuranosidase activity. Such a homologue can be selected and produced by those skilled in the art without undue experiments by referring to the sequence of SEQ ID No. 1 or 3.

[0015] The β -fructofuranosidase having the amino acid sequence of SEQ ID Nos. 1 and 3 according to the present invention has high fructosyltransferase activity and efficiently produces fructooligosaccharides. Specifically, when a sucrose solution at a concentration of 30 wt% or more is used as a substrate for reaction, the fructosyltransferase activity of β -fructofuranosidase having the amino acid sequence of SEQ ID No. 1 is at least 4 times higher, and the fructosyltransferase activity of β -fructofuranosidase having the amino acid sequence of SEQ ID No. 3 is at least 7 times higher than hydrolytic activity. Furthermore, 50% or more of sucrose is converted to fructooligosaccharides in both cases.

 β -fructofuranosidase gene

[0016] The novel gene encoding β -fructofuranosidase according to the present invention comprises a DNA sequence encoding the amino acid sequence of SEQ ID Nos. 1 and 3 or a homologue thereof.

[0017] Generally, a nucleotide sequence which encodes the amino acid sequence of a given protein can be easily determined from the reference chart known as "codon table". A variety of nucleotide sequence are available from those encoding the amino acid sequence of SEQ ID No. 1 or 3. Therefore, the term "a nucleotide sequence encoding the amino acid sequence of SEQ ID No. 1 or 3" refers to the meaning including the nucleotide sequence of SEQ ID No. 2 or 4, as well as nucleotide sequences which consist of the same codons as above allowing for degeneracy and encode the amino acid sequence of SEQ ID No. 1 or 3.

[0018] A preferred embodiment of the present invention provides, as a preferred example of the novel gene according to the present invention, a DNA fragment comprising the nucleotide sequence of SEQ ID No. 2 or 4.

[0019] As described above, the present invention encompasses a homologue of the amino acid sequence of SEQ ID No. 1 or 3. Therefore, the DNA fragment according to the present invention involves a nucleotide sequence which encodes such a homologue.

[0020] As the nucleotide sequence of the DNA fragment according to the present invention is determined, the DNA fragment may be obtained according to the procedure for the synthesis of a nucleic acid.

[0021] This sequence can also be obtained from Penicillium roqueforti or Scopulariopsis brevicaulis, preferably

EP 1 022 332 A1

Penicillium roqueforti IAM7254 or Scopulariopsis brevicaulis IFO4843, according to the procedure of genetic engineering.

Expression of β -fructofuranosidase Gene

[0022] The β -fructofuranosidase according to the present invention can be produced in a host cell which has been transformed by a DNA fragment encoding the enzyme. More specifically, a DNA fragment encoding the β -fructofuranosidase according to the present invention is introduced in a host cell in the form of a DNA molecule which is replicatable in the host cell and can express the above gene, particularly an expression vector, in order to transform the host cell. Then, the obtained transformant is cultivated.

[0023] Therefore, the present invention provides a DNA molecule which comprises a gene encoding the β -fructofuranosidase according to the present invention, particularly an expression vector. This DNA molecule is obtained by introducing a DNA fragment encoding the β -fructofuranosidase according to the present invention in a vector molecule. According to a preferred embodiment of the present invention, the vector is a plasmid.

[0024] The DNA molecule according to the present invention may be prepared by the standard technique of genetic engineering.

[0025] The vector applicable in the present invention can be selected as appropriate from viruses, plasmids, cosmid vectors, etc., considering the type of the host cell used. For example, a bacteriophage in the λ phage group or a plasmid in the pBR or pUC group may be used for E. coli host cells, a plasmid in the pUB group for Bacillus subtilis, and a vector in the YEp or YCp group for yeast.

[0026] It is preferable that the plasmid contain a selectable marker to ensure the selection of the obtained transformance, such as a drug-resistance marker or marker gene complementing an auxotrophic mutation. Preferred example of marker genes include ampicillin-resistance gene, kanamycin-resistance gene, and tetracycline-resistance gene for bacterium host cells; N-(5'-phosphoribosyl)-anthranilate isomerase gene (TRP1), orotidine-5'-phosphate decarboxylase gene (URA3), and β -isopropylmalate dehydrogenase gene (LEU2) for yeast; and hygromycin-resistance gene (hph), bialaphos-resistance gene (bar), and nitrate reductase gene (niaD) for mold.

[0027] It is also preferable that the DNA molecule for use as an expression vector according to the present invention contain nucleotide sequence necessary for the expression of the β -fructofuranosidase gene, including transcription and translation control signals, such as a promoter, a transcription initiation signal, a ribosome binding site, a translation termination signal, and a transcription termination signal.

[0028] Examples of preferred promoters include, in addition to the promoter on the inserted fragment which is able to function in the host, promoters such as those of lactose operon (lac), and tryptophan operon (trp) for E. coli; promoters such as those of alcohol dehydrogenase gene (ADH), acid phosphatase gene (PHO), galactose regulated gene (GAL), and glyceraldehyde-3-phosphate dehydrogenase gene (GPD) for yeast; and promoters such as those of α -amylase gene (amy) and cellobiohydrolase I gene (CBHI) for mold.

[0029] When the host cell is Bacillus subtilis, yeast or mold, it is also advantageous to use a secretion vector to allow it to extracellularly secrete the produced recombinant β -fructofuranosidase. Any host cell with an established host-vector system may be used, preferably yeast, mold, etc. It is preferable also to use the mold fungus having no β -fructofuranosidase activity described in PCT/JP97/00757.

[0030] A novel recombinant enzyme produced by the transformant described above is obtained by the following procedure: first, the host cell described above is cultivated under suitable conditions to obtain the supernatant or cell bodies from the resultant culture, using a known technique such as centrifugation; cell bodies should be further suspended in a suitable buffer solution, then homogenized by freeze-and-thaw, ultrasonic treatment, or mortar, followed by centrifugation or filtration to separate a cell body extract containing the novel recombinant enzyme.

[0031] The enzyme can be purified by combining the standard techniques for separation and purification. Examples of such as techniques include processes such as heat treatment, which rely on the difference in thermal resistance; processes such as salt sedimentation and solvent sedimentation, which rely on the difference in solubility; processes such as dialysis, ultrafiltration and gel filtration, and SDS-polyacrylamide gel electrophoresis, which rely on the difference in molecular weight; processes such as ion exchange chromatography, which rely on the difference in electric charge; processes such as affinity chromatography, which rely on specific affinity; processes such as hydrophobic chromatography and reversed-phase partition chromatography, which rely on the difference in hydrophobicity; and processes such as isoelectric focusing, which rely on the difference in isoelectric point.

Production of fructooligosaccharides using the β -fructofuranosidase

[0032] The present invention further provides a process for producing fructooligosaccharide using the recombinant host or recombinant β -fructofuranosidase described above.

[0033] In the process for producing fructooligosaccharides according to the present invention, the recombinant host

EP 1 022 332 A1

or recombinant β -fructofuranosidase described above is brought into contact with sucrose.

[0034] The mode and conditions where the recombinant host or recombinant β -fructofuranosidase according to the present invention comes in contact with sucrose are not limited in any way provided that the novel recombinant enzyme is able to act on sucrose. A preferred embodiment for contact in solution is as follows: The sucrose concentration may be selected as appropriate in the range where sucrose can be dissolved. However, considering the conditions such as the specific activity of the enzyme and reaction temperature, the concentration should generally fall in the range of 5% to 80%, preferably 30% to 70%. The temperature and pH for the reaction of sucrose by the enzyme should preferably be optimized for the characteristics of the novel recombinant enzyme. Therefore, the reasonable conditions are about 30°C to 80°C, pH 4 to 10, preferably 40°C to 70°C, pH 5 to 7.

[0035] The degree of purification of the novel recombinant enzyme may be selected as appropriate. The enzyme may be used either as unpurified in the form of supernatant from a transformant culture or cell body homogenate, as purified after processed in various purification steps, or as isolated after processed by various purification means.

[0036] Furthermore, the enzyme may be brought into contact with sucrose as fixed on a carrier using the standard technique.

[0037] The fructooligosaccharides thus produced are purified from the resulting solution according to known procedures. For example, the solution may be heated to inactivate the enzyme, decolorized using activated carbon, then desalted using ion exchange resin.

Examples

Example 1: Determination of translation region of β -fructofuranosidase gene from Penicillium roqueforti IAM7254

[0038] A DNA fragment of about 2 kbp containing the β -fructofuranosidase gene from Aspergillus niger was amplified by PCR using a chromosomal DNA prepared from Aspergillus niger ATCC20611 according to the standard procedure as a template and synthetic DNAs of SEQ ID Nos. 5 and 6 as primers. This DNA fragment was fractionated by agarose gel electrophoresis, extracted according to the standard procedure, purified, and then dissolved in sterilized water to 0.1 $\mu\text{g}/\mu\text{l}$ to prepare a DNA sample for probe.

[0039] In the next step, a chromosomal DNA from Penicillium roqueforti IAM7254 was prepared, about 20 μg of the chromosomal DNA was digested completely with EcoRI, followed by agarose gel electrophoresis to recover about 4 kbp DNA fragments.

[0040] The recovered DNA fragments of about 4 kbp (about 0.5 μg) were ligated with 1 μg of $\lambda\text{gt}10$ vector which had been digested with EcoRI and treated with phosphatase, packaged using an *in vitro* packaging kit, GIGAPACK II Gold (Stratagene L.L.C.), and then introduced in E. coli NM514, to prepare a library.

[0041] A probe was prepared from DNA sample for probe above described. As a result of plaque hybridization using ECL Direct DNA/RNA Labelling & Detection System (Amersham International), 4 clones turned out positive in about 25,000 plaques. These positive clones were purified by a second screening to prepare phage DNA, which was then analyzed using restriction enzymes. The result showed that all the clones had an identical EcoRI fragment of about 4 kbp.

[0042] The EcoRI fragments of about 4 kbp were subdivided into a small fragment to select the desired DNA region using restriction enzymes, then subcloned to plasmid vector pUC118 or pUC119. The plasmid DNA was obtained from the subclone according to the standard procedure and sequenced using ALFred DNA Sequencer (Pharmacia) as shown in SEQ ID No.7.

[0043] The sequence consisting 50 bases from 1695 to 1744 in this sequence was identified as an intron because it showed a typical intron structure of filamentous fungi. As a result, the sequence of SEQ ID No. 2 as a sequence encoding protein was obtained by deleting the intron from the sequence of SEQ ID No. 7. The encoded amino acid sequence was shown in SEQ ID No. 1.

Example 2: Expression of β -fructofuranosidase gene from Penicillium roqueforti IAM7254 in Saccharomyces cerevisiae

[0044] Plasmid pYPEN01 and pYPEN02 for expressing the β -fructofuranosidase gene from Penicillium roqueforti were prepared as follows (Figure 1A, B, C and D).

[0045] pYPR2831 (H. Horiuchi et al., Agric. Biol. Chem., 54, 1771-1779, 1990) was digested with EcoRI and Sall, and then its terminals were blunted with T4 DNA polymerase. The obtained fragment was ligated with BamHI linker (5'-CGGATCCG-3'), digested with BamHI, followed by self-ligation to obtain vector pY2831 for expression in yeast.

[0046] Next, single-stranded DNA was prepared from the plasmid pPRS01 obtained by inserting an about 4 kbp EcoRI DNA fragment containing the β -fructofuranosidase gene prepared in Example 1 into plasmid pUC118. Using the single-stranded DNA as a template and a synthetic DNA of SEQ ID No. 8 as a primer, the translated region of the β -fructofuranosidase gene was subjected to site-specific mutagenesis to disrupt the BamHI site without changing the

EP 1 022 332 A1

encoded amino acid sequence (pPRS02).

[0047] A part of the translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using plasmid pPRS02 as a template and synthetic DNAs of SEQ ID Nos. 9 and 10 as primers, and inserted into the BamHI site of plasmid pY2831 to prepare pYPEN01. Thus, plasmid pYPEN01 is designed to produce an enzyme protein comprising an amino acid sequence from 1 to 565 in the amino acid sequence of SEQ ID No. 1, which is a mature β -fructofuranosidase following secretion signal sequence.

[0048] Further, a DNA fragment containing the translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using plasmid pPRS02 as a template and synthetic DNAs of SEQ ID Nos. 9 and 11 as primers, and inserted into the BamHI site of plasmid pUC118 to prepare plasmid pPRS03. A single-stranded DNA was prepared from plasmid pPRS03. As a result of site-specific mutagenesis using this as a template and a synthetic DNA of SEQ ID No. 12 as a primer, the intron sequence was removed (pPRS04). The translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment from plasmid pPRS04, and inserted into the BamHI site of plasmid pY2831 to prepare plasmid pYPEN02. Thus, plasmid pYPEN02 is designed to produce an enzyme protein comprising an amino acid sequence of SEQ ID No. 1, which is a mature β -fructofuranosidase following secretion signal sequence.

[0049] Plasmids pYPEN01 and pYPEN02 were introduced into *Saccharomyces cerevisiae* MS-161 (Suc⁺, ura3, trp1) by the lithium-acetate method (Ito, H. et al., J. Bacteriol., 153, 163-168, 1983) to obtain transformants. The transformants were cultivated in an SD-Ura medium (0.67% yeast nitrogen base (Difco), 2% glucose and 50 μ g/ml uracil) at 30°C overnight. The culture was seeded in a production medium (0.67% yeast nitrogen base Difco, 2% glucose, 2% casamino acid and 50 μ g/ml uracil) at a final concentration of 1% and cultivated at 30°C for 2 days. The culture supernatant was analyzed for β -fructofuranosidase activity, in units, i.e., the quantity of free glucose (μ mol) released in 1 minute in 10 wt% sucrose solution, pH 5.5, at 40°C for 60 minutes. As a result, the transformant with plasmid pYPEN01 exhibited 4×10^{-4} units/ml or less of activity while the transformant with plasmid pYPEN02 exhibited 0.38 units/ml of activity.

Example 3: Determination of the translated region of β -fructofuranosidase gene from *Scopulariopsis brevicaulis* IFO4843

[0050] The chromosomal DNA was prepared from *Scopulariopsis brevicaulis* IFO4843. About 20 μ g of a chromosomal DNA sample was completely digested with EcoRI, and electrophoresed through an agarose gel to recover an about 10 kbp DNA fragment.

[0051] The recovered DNA fragment of about 10 kbp (about 0.5 μ g) were ligated with 1 μ g of λ DASHII vector digested with HindIII and EcoRI, and packaged using an *in vitro* packaging kit, GIGAPACK II Gold (Stratagene L.L.C.), then introduced in *E. coli* XL1-Blue MRA (P2), to prepare a library.

[0052] As a result of plaque hybridization using ECL Direct DNA/RNA Labelling & Detection System (Amersham International), with the about 2 kbp DNA fragment used in Example 1 as a probe, 3 clones turned out positive in about 15,000 plaques. These positive clones were purified by a second screening to prepare phage DNA, which was then analyzed using restriction enzymes. The result showed that all the clones had an identical EcoRI fragment of about 10 kbp.

[0053] These EcoRI fragments of about 10 kbp were subdivided into a small fragment to select the desired DNA region using restriction enzymes, then subcloned to plasmid vector pUC118 or pUC119. The plasmid DNA was obtained from the subclone according to the standard procedure and sequenced using ALFred DNA Sequencer (Pharmacia) as shown in SEQ ID No.13.

[0054] The sequence comprising 55 bases from 1722 to 1776 in this sequence was identified as an intron because it showed a typical intron structure of filamentous fungi. As a result, the sequence of SEQ ID No. 4 as a sequence encoding protein was obtained by deleting the intron from the sequence of SEQ ID No. 13. The encoded amino acid sequence was shown SEQ ID No. 3.

Example 4: Expression of β -fructofuranosidase gene from *Scopulariopsis brevicaulis* IFO4843 in *Saccharomyces cerevisiae*

[0055] Plasmid pYSCOP01 and pYSCOP02 for expressing the β -fructofuranosidase gene from *Scopulariopsis brevicaulis* were prepared as follows (Figure 2A and B).

[0056] A part of the translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using about 10 kbp EcoRI DNA fragment prepared in Example 3 containing the β -fructofuranosidase gene as a template and synthetic DNAs of SEQ ID Nos. 14 and 15 as primers, and inserted into the BamHI site of plasmid pY2831 to prepare pYSCOP01. Thus, plasmid pYSCOP01 is designed to produce an enzyme protein comprising an amino acid sequence from 1 to 574 in the amino acid sequence of SEQ ID No. 3, which is a mature β -fructofuranosi-

EP 1 022 332 A1

dase following secretion signal sequence.

[0057] Next, a DNA fragment containing the translated region of the β -fructofuranosidase gene was prepared as an about 1.9 kbp BamHI fragment by PCR using an about 10 kbp EcoRI fragment containing the β -fructofuranosidase gene as a template and synthetic DNAs of SEQ ID Nos. 14 and 16 as primers, and inserted into the BamHI site of plasmid pUC118 to prepare plasmid pSCB01. A single-stranded DNA was prepared from plasmid pSCB01. As a result of site-specific mutagenesis using this as a template and the synthetic DNA of SEQ ID No. 17 as a primer, the intron sequence was removed (pSCB02). The translated region of the β -fructofuranosidase gene was prepared as an about 1.9 kbp BamHI fragment from plasmid pSCB02, and inserted into the BamHI site of plasmid pY2831 to prepare plasmid pYSCOP02. Thus, plasmid pYSCOP02 is designed to produce an enzyme protein comprising an amino acid sequence of SEQ ID No. 3, which is a mature β -fructofuranosidase following secretion signal sequence.

[0058] Plasmids pYSCOP01 and pYSCOP02 were introduced into *Saccharomyces cerevisiae* MS-161 (Suc⁺, ura3, trp1) by the lithium-acetate method to obtain transformants. The transformants were cultivated in an SD-Ura medium at 30°C overnight. The culture was seeded a production medium at a final concentration of 1% and cultivated at 30°C for 2 days. The culture supernatant was analyzed for β -fructofuranosidase activity in the same manner as described in Example 2. As a result, the transformant with plasmid pYSCOP01 exhibited 4×10^{-4} units/ml or less of activity, while the transformant with plasmid pYSCOP02 exhibited 6.5×10^{-3} units/ml of activity.

EP 1 022 332 A1

SEQUENCE LISTING

5

<110> MEIJI SEIKA KAISHA, LTD.

10

<120> Beta-fructofuranosidase and its gene

<130> 116875-475

15

<140>

<141> 1998-9-10

20

<150> JP/245154/1997

<151> 1997-9-10

25

<160> 17

30

<170> PatentIn Ver. 2.0

35

<210> 1

<211> 603

<212> PRT

40

<213> Penicillium roqueforti IAM7254

<220>

<221> mat peptide

45

<222> (1)... (603)

50

<400> 1

Val Asp Phe His Thr Pro Ile Asp Tyr Asn Ser Ala Pro Pro Asn Leu

1

5

10

15

55

EP 1 022 332 A1

Ser Thr Leu Ala Asn Ala Ser Leu Phe Lys Thr Trp Arg Pro Arg Ala
 20 25 30
 5 His Leu Leu Pro Pro Ser Gly Asn Ile Gly Asp Pro Cys Gly His Tyr
 35 40 45
 10 Thr Asp Pro Lys Thr Gly Leu Phe His Val Gly Trp Leu Tyr Ser Gly
 50 55 60
 Ile Ser Gly Ala Thr Thr Asp Asp Leu Val Thr Tyr Lys Asp Leu Asn
 15 65 70 75 80
 Pro Asp Gly Ala Pro Ser Ile Val Ala Gly Gly Lys Asn Asp Pro Leu
 85 90 95
 20 Ser Val Phe Asp Gly Ser Val Ile Pro Ser Gly Ile Asp Gly Met Pro
 100 105 110
 Thr Leu Leu Tyr Thr Ser Val Ser Tyr Leu Pro Ile His Trp Ser Ile
 25 115 120 125
 Pro Tyr Thr Arg Gly Ser Glu Thr Gln Ser Leu Ala Val Ser Tyr Asp
 130 135 140
 30 Gly Gly His Asn Phe Thr Lys Leu Asn Gln Gly Pro Val Ile Pro Thr
 145 150 155 160
 Pro Pro Phe Ala Leu Asn Val Thr Ala Phe Arg Asp Pro Tyr Val Phe
 35 165 170 175
 Gln Ser Pro Ile Leu Asp Lys Ser Val Asn Ser Thr Gln Gly Thr Trp
 40 180 185 190
 Tyr Val Ala Ile Ser Gly Gly Val His Gly Val Gly Pro Cys Gln Phe
 195 200 205
 45 Leu Tyr Arg Gln Asn Asp Ala Asp Phe Gln Tyr Trp Glu Tyr Leu Gly
 210 215 220
 Gln Trp Trp Lys Glu Pro Leu Asn Thr Thr Trp Gly Lys Gly Asp Trp
 50 225 230 235 240
 Ala Gly Gly Trp Gly Phe Asn Phe Glu Val Gly Asn Val Phe Ser Leu
 55

EP 1 022 332 A1

	245	250	255
5	Asn Ala Glu Gly Tyr Ser Glu Asp Gly Glu Ile Phe Ile Thr Leu Gly		
	260	265	270
	Ala Glu Gly Ser Gly Leu Pro Ile Val Pro Gln Val Ser Ser Ile Arg		
10	275	280	285
	Asp Met Leu Trp Val Thr Gly Asn Val Thr Asn Asp Gly Ser Val Thr		
	290	295	300
15	Phe Lys Pro Thr Met Ala Gly Val Leu Asp Trp Gly Val Ser Ala Tyr		
	305	310	315
	Ala Ala Ala Gly Lys Ile Leu Pro Ala Ser Ser Gln Ala Ser Thr Lys		
20	325	330	335
	Ser Gly Ala Pro Asp Arg Phe Ile Ser Tyr Val Trp Leu Thr Gly Asp		
	340	345	350
25	Leu Phe Glu Gln Val Lys Gly Phe Pro Thr Ala Gln Gln Asn Trp Thr		
	355	360	365
	Gly Ala Leu Leu Leu Pro Arg Glu Leu Asn Val Arg Thr Ile Ser Asn		
30	370	375	380
	Val Val Asp Asn Glu Leu Ser Arg Glu Ser Leu Thr Ser Trp Arg Val		
	385	390	395
35	Ala Arg Glu Asp Ser Gly Gln Ile Asp Leu Glu Thr Met Gly Ile Ser		
	405	410	415
	Ile Ser Arg Glu Thr Tyr Ser Ala Leu Thr Ser Gly Ser Ser Phe Val		
40	420	425	430
	Glu Ser Gly Lys Thr Leu Ser Asn Ala Gly Ala Val Pro Phe Asn Thr		
	435	440	445
45	Ser Pro Ser Ser Lys Phe Phe Val Leu Thr Ala Asn Ile Ser Phe Pro		
	450	455	460
50	Thr Ser Ala Arg Asp Ser Gly Ile Gln Ala Gly Phe Gln Val Leu Ser		
	465	470	475
55			

EP 1 022 332 A1

Ser Ser Leu Glu Ser Thr Thr Ile Tyr Tyr Gln Phe Ser Asn Glu Ser

485

490

495

Ile Ile Val Asp Arg Ser Asn Thr Ser Ala Ala Ala Arg Thr Thr Ala

500

505

510

Gly Ile Leu Ser Asp Asn Glu Ala Gly Arg Leu Arg Leu Phe Asp Val

515

520

525

Leu Arg Asn Gly Lys Glu Gln Val Glu Thr Leu Glu Leu Thr Ile Val

530

535

540

Val Asp Asn Ser Val Leu Glu Val Tyr Ala Asn Gly Arg Phe Ala Leu

545

550

555

560

Gly Thr Trp Ala Arg Ser Trp Tyr Ala Asn Ser Thr Lys Ile Asn Phe

565

570

575

Phe His Asn Gly Val Gly Glu Ala Thr Phe Glu Asp Val Thr Val Phe

580

585

590

Glu Gly Leu Tyr Asp Ala Trp Pro Gln Arg Lys

595

600

<210> 2

<211> 1809

<212> DNA

<213> Penicillium roqueforti IAM7254

<220>

<221> mat peptide

<222> (1)... (1809)

<400> 2

gttgatttcc ataccccgat tgactataac tgggtccgc caaaccttcc taccctggca 60

aacgcatttc ttttcaagac atggagacc agageccatc ttctccctcc atcigggaac 120

ataggcgacc cgtgcgggca clataccgat cccaagactg gtctcttcca cgtgggttgg 180

EP 1 022 332 A1

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

ctttacagtg ggatttcggg agcgacaacc gacgatctcg ttacctataa agacctcaat 240
 cccgatggag ccccgicaat tgitgcagga ggaagaacg accctcttct tctcttcgat 300
 ggctcggica ttccaagcgg tatagacggc atgccaactc ttctgtatag ctctgtatca 360
 tacctcccaa tccactggtc catccctac acccggggaa gcgagacaca atccctggcc 420
 gtttccatg acggiggtca caacttcacc aagctcaacc aagggcccgat gatccctacg 480
 cctccgittg ctctcaatgt caccgcttct cgtgacccct acgttttcca aagcccaatt 540
 ctggacaaat ctgtcaatag tacccaagga acatgggatg tgcctatctc tggcgggtgc 600
 cacgggtgctg gaccttgctc gtctctctac cgtcagaacg acgcagattt tcaatatagg 660
 gaatatctcg ggcaatggig gaaggagccc cttaatacca ctgggggaaa gggtgactgg 720
 gccgggggtt ggggcctcaa ctltgagggt ggcaacgtct ttagtctgaa tgcagagggg 780
 tatagtgag acggcgagat attcataacc ctgggtgctg aggggtcggg acttcccatc 840
 gtctctcaag tctctctat tgcgatatg ctgtgggtga ccggcaatgt cacaaatgac 900
 ggctctgtca ctltcaagcc aacctggcg gggtgcttg actggggcgt gtccgcatat 960
 gctgctgcag gcaagatctt gccggccagc tctcaggcat ccacaaagag cggtgcccc 1020
 gatcggitca ttctctatgt ctggctcact ggagatctat tgcagcaagt gaaaggattc 1080
 cctaccgctc aacaaaactg gaccggggcc ctcttactgc ccgagagct gaatgtccgc 1140
 actatctcta acgtgggtga taacgaactt tgcgtgagt ccttgacatc gtggcgcggtg 1200
 gcccgcaag actctggca gatcgacctt gaaacaatgg gaatctcaat tccaggzag 1260
 acttacagcg ctctcctac cggctcatct ttgtcgagt ctggtaaac ctctcgzat 1320
 gctggagcag tgccttcaa tacctcacc tcaagcaagt tcttcgtgct gacagcaaat 1380
 atatcttcc cgacctctgc ccgtgactct ggcatccagg ctggtttcca ggttttatcc 1440
 tctagtttg agtctacaac tatctactac caattctcca acgagtcct catcgtcgac 1500
 cgcagcaaca cgagtgctgc ggcgagaaca actgctggga tcttcagtga taacgaggcg 1560
 ggacgtctgc gcccttctga cgtgttgca aatggaaaag aacaggtiga aactttggag 1620
 ctactatcg tggtgataa tagtgtactg gaagtatatg ccaatggacg ctttgctcta 1680
 ggcactlggg ctgggtcttg gtacgccaac tgcactaaa ttaacttctt ccataacggc 1740
 gtgggagaag cgacattcga agatgtgacg gtctttgaag gactgtatga tgcctggcca 1800
 caaaggaag 1809

EP 1 022 332 A1

<210> 3

<211> 613

<212> PRT

<213> Scopulariopsis brevicaulis IF04843

<220>

<221> mat peptide

<222> (1)... (613)

<400> 3

Gln Pro Thr Ser Leu Ser Ile Asp Asn Ser Thr Tyr Pro Ser Ile Asp

1 5 10 15

Tyr Asn Ser Ala Pro Pro Asn Leu Ser Thr Leu Ala Asn Asn Ser Leu

20 25 30

Phe Glu Thr Trp Arg Pro Arg Ala His Val Leu Pro Pro Gln Asn Gln

35 40 45

Ile Gly Asp Pro Cys Met His Tyr Thr Asp Pro Glu Thr Gly Ile Phe

50 55 60

His Val Gly Trp Leu Tyr Asn Gly Asn Gly Ala Ser Gly Ala Thr Thr

65 70 75 80

Glu Asp Leu Val Thr Tyr Gln Asp Leu Asn Pro Asp Gly Ala Gln Met

85 90 95

Ile Leu Pro Gly Gly Val Asn Asp Pro Ile Ala Val Phe Asp Gly Ala

100 105 110

Val Ile Pro Ser Gly Ile Asp Gly Lys Pro Thr Met Met Tyr Thr Ser

115 120 125

Val Ser Tyr Met Pro Ile Ser Trp Ser Ile Ala Tyr Thr Arg Gly Ser

130 135 140

Glu Thr His Ser Leu Ala Val Ser Ser Asp Gly Gly Lys Asn Phe Thr

145 150 155 160

EP 1 022 332 A1

Lys Leu Val Gln Gly Pro Val Ile Pro Ser Pro Pro Phe Gly Ala Asn

165

170

175

Val Thr Ser Trp Arg Asp Pro Phe Leu Phe Gln Asn Pro Gln Phe Asp

180

185

190

Ser Leu Leu Glu Ser Glu Asn Gly Thr Trp Tyr Thr Val Ile Ser Gly

195

200

205

Gly Ile His Gly Asp Gly Pro Ser Ala Phe Leu Tyr Arg Gln His Asp

210

215

220

Pro Asp Phe Gln Tyr Trp Glu Tyr Leu Gly Pro Trp Trp Asn Glu Glu

225

230

235

240

Gly Asn Ser Thr Trp Gly Ser Gly Asp Trp Ala Gly Arg Trp Gly Tyr

245

250

255

Asn Phe Glu Val Ile Asn Ile Val Gly Leu Asp Asp Asp Gly Tyr Asn

260

265

270

Pro Asp Gly Glu Ile Phe Ala Thr Val Gly Thr Glu Trp Ser Phe Asp

275

280

285

Pro Ile Lys Pro Gln Ala Ser Asp Asn Arg Glu Met Leu Trp Ala Ala

290

295

300

Gly Asn Met Thr Leu Glu Asp Gly Asp Ile Lys Phe Thr Pro Ser Met

305

310

315

320

Ala Gly Tyr Leu Asp Trp Gly Leu Ser Ala Tyr Ala Ala Ala Gly Lys

325

330

335

Glu Leu Pro Ala Ser Ser Lys Pro Ser Gln Lys Ser Gly Ala Pro Asp

340

345

350

Arg Phe Val Ser Tyr Leu Trp Leu Thr Gly Asp Tyr Phe Glu Gly His

355

360

365

Asp Phe Pro Thr Pro Gln Gln Asn Trp Thr Gly Ser Leu Leu Leu Pro

370

375

380

Arg Glu Leu Ser Val Gly Thr Ile Pro Asn Val Val Asp Asn Glu Leu

EP 1 022 332 A1

385 390 395 400
Ala Arg Glu Thr Gly Ser Trp Arg Val Gly Thr Asn Asp Thr Gly Val
405 410 415
Leu Glu Leu Val Thr Leu Lys Gln Glu Ile Ala Arg Glu Thr Leu Ala
420 425 430
Glu Met Thr Ser Gly Asn Ser Phe Thr Glu Ala Ser Arg Asn Val Ser
435 440 445
Ser Pro Gly Ser Thr Ala Phe Gln Gln Ser Leu Asp Ser Lys Phe Phe
450 455 460
Val Leu Thr Ala Ser Leu Ser Phe Pro Ser Ser Ala Arg Asp Ser Asp
465 470 475 480
Leu Lys Ala Gly Phe Glu Ile Leu Ser Ser Glu Phe Glu Ser Thr Thr
485 490 495
Val Tyr Tyr Gln Phe Ser Asn Glu Ser Ile Ile Ile Asp Arg Ser Asn
500 505 510
Ser Ser Ala Ala Ala Leu Thr Thr Asp Gly Ile Asp Thr Arg Asn Glu
515 520 525
Phe Gly Lys Met Arg Leu Phe Asp Val Val Glu Gly Asp Gln Glu Arg
530 535 540
Ile Glu Thr Leu Asp Leu Thr Ile Val Val Asp Asn Ser Ile Val Glu
545 550 555 560
Val His Ala Asn Gly Arg Phe Ala Leu Ser Thr Trp Val Arg Ser Trp
565 570 575
Tyr Glu Ser Ser Lys Asp Ile Lys Phe Phe His Asp Gly Asp Ser Thr
580 585 590
Val Gln Phe Ser Asn Ile Thr Val Tyr Glu Gly Leu Phe Asp Ala Trp
595 600 605
Pro Glu Arg Ala Arg
610

EP 1 022 332 A1

<210> 4

<211> 1839

<212> DNA

<213> Scopulariopsis brevicaulis IF04843

<220>

<221> mat peptide

<222> (1)... (1839)

<400> 4

20	caacctacgt ctcgtcaat cgacaattc acgtatcctt ctatcgacta caactccgcc	60
	cctccaaacc tctcgactct tgccaacaac agcctcttcg agacatggag gccgaggcca	120
	cacgtccttc cgccccagaa ccagatcggc gatccgltga tgcactacac cgaccccgag	180
25	acaggaatct tccacgtcgg ctggctgtac aacggcaatg gcgcttcgg cgccacgacc	240
	gaggatctcg tcacctatca ggatctcaac cccgacggag cgcagatgat ccttcgggt	300
	ggigtgaatg accccattgc tgtctttgac ggcgcggtta ttcccagtgg cattgatggg	360
30	aaaccaccca tgatgtatac ctcggtgtca tacatgccca tctcctggag catcgcttac	420
	accaggggaa gcgagaccca ctctctcgca gtgtcgtccg acggcggtta gaacttcacc	480
	aagctgggtc agggccccgi cattccttcg cctcccttcg gcgccaacgt gaccagctgg	540
35	cgtgaccctt tctgtttca aaacccccag ttcgactctc tctcgaaag cgagaacggc	600
	acgtggtaca ccgttatctc tggtagcatc cacggtgacg gccccccgc gtctctctac	660
40	cgtcagcacg accccgactt ccagtactgg gagtacctg gaccgtgggt gaacgaggaa	720
	gggaactcga cctggggcag cgtgactgg gctggccggt ggggctacaa cttcgaggtc	780
	atcaacattg tcggtcttga cgatgatggc tacaaccccg acggtgaaat ctttgccacg	840
45	gtaggtaccg aatggtcgtt tgaccccatc aaaccgcagg cctcggacaa caggagatg	900
	ctctgggccc cgggcaacat gactctcgag gacggcgata tcaagttcac gccaaagcatg	960
	gcgggctacc tcgactgggg tctatcgggc tatgccgccc ctggcaagga gctgcccgt	1020
50	tcttcaaagc cttcgcagaa gagcgggtgc cggaccggt tcgtgtcgta cctgtggctc	1080
	accggtgact acttcgaggg ccacgacttc cccaccccg cgcagaattg gaccggctcg	1140

EP 1 022 332 A1

5 cttttgcttc cgcgtgagct gagcgtcggg acgattccca acgttgtcga caacgagctt 1200
gctcgcgaga cgggctcttg gagggttggc accaaccgaca ctggcgtgct tgagctggtc 1260
actctgaagc aggagattgc tcgcgagacg ctggctgaaa tgaccagcgg caactccttc 1320
10 accgaggcga gcaggaatgt cagctcgccc ggatctaccg ccttccagca gtccctggat 1380
tccaagttct tcgtcttgac cgcctcgctc tcttccctt cgtcggctcg cgactccgac 1440
ctcaaggctg gtttcgagat cctgtcgtcc gagtttgagt cgaccacggt ctactaccag 1500
ttttccaacg agtccatcat cattgaccgg agcaactcga gtgcgtccgc cttgactacc 1560
15 gatggaatcg acacccgcaa cgagtttggc aagatgcgcc tgtttgatgt tctcgagggt 1620
gaccaggagc gtaicgagac gctcgatctc actatgttgg ttgataactc gatcgttgag 1680
gttcatgcca acgggcgatt cgtctgagc acttgggttc gttcgttgta cgagtcgtcc 1740
20 aaggacatca agttcttcca cgalggcgac agcacggttc agttctcgaa catcacgctc 1800
tacgagggac tgtttgacgc ctggccggag cgggccagg 1839

25 <210> 5

<211> 20

30 <212> DNA

<213> Artificial Sequence

35 <400> 5

caatgaagct caccactacc

40 <210> 6

<211> 20

45 <212> DNA

<213> Artificial Sequence

50 <400> 6

atcccggtca atttctctcc

EP 1 022 332 A1

<210> 7

<211> 1809

<212> DNA

<213> *Penicillium roqueforti* IAM7254

<220>

<221> CDS

<222> (1)... (1694)

<221> intron

<222> (1695)... (1744)

<221> CDS

<222> (1745)... (1859)

<400> 1

gttagatttcc atacccegat tgactataac tgggtccgc caaaccttcc taccctggca 60
 aacgcattcc ttttcaagac atggagacc agagcccatc ttctccctcc atctgggaac 120
 ataggcgacc cgtgcgggca ctataccgat cccaagactg gtctcttcca cgtgggttgg 180
 ctttacagtg ggatttcggg agcgacaacc gacgatctcg ttacctataa agacctcaat 240
 cccgatggag ccccgicaat tgttgcagga ggaaagaacg accctctttc tgtcttcgat 300
 ggctcggta tccaagcgg tatagacggc atgccaactc ttctgtatca ctctgtatca 360
 tacttccaa tccactggc calccctac acccggggaa gcgagacaca atccttggcc 420
 gtttccatg acggtggta caacttcacc aagctcaacc aagggcccgat gatccctacg 480
 cctccgtttg ctctcaatgt caccgcttc cgtgaccctc acgttttcca aagcccaatt 540
 ctggacaaat ctgtcaatag tacccaagga acatggatg tgcctatc tggcgggtgc 600
 cacgggtgc gaccttgta gtctctctac cgtcagaacg acgcagattt tcaatatgg 660
 gaatatctcg ggcaatggg gaaggagccc cttaatacca ctgggggaaa gggtagctgg 720
 gccgggggtt ggggcctcaa cttgaggtt ggcaacgtct ttagtctgaa tgcagagggg 780
 tatagtgaag acggcgagat attcataacc ctccgtgctg agggttcggg acttcccatc 840
 gtacctcaag tctctctat tgcgataatg ctgtgggtga ccggcaatgt cacaaatgac 900

EP 1 022 332 A1

5
10
15
20
25
30
35
40
45
50
55

ggctctgtca ctttcaagcc aacctggcg ggtgtgttg actggggcgt gtcggcata 960
gtgtgtcag gcaagatctt gccggccagc tctcaggcat ccacaaagag cggtgcccc 1020
gatcggttca tttcclatgt ctggctcact ggagatctat tcgagcaagt gaaaggattc 1080
cctaccgctc aacaaaactg gaccggggcc ctcttactgc cgcgagagct gaatgtccgc 1140
actatctcta acgtgggtga taacgaactt tcgcgtgagt ccttgacatc gtggcgcgtg 1200
gcccgcgaag actctgttca gatcgacctt gaaacaatgg gaatctcaat ttccaggag 1260
acttacagcg ctctcacatc cggtcactt ttgtctgagt ctggtaaaac gttgtcgaat 1320
gctggagcag tgcccttcaa tacctcacc ccaagcaagt tcttcgtgtc gacagcaat 1380
atatctttcc cgacctctgc ccgtgactct ggcatccagg ctggtttcca ggttttatcc 1440
tctagtcttg agtctacaac tatctactac caattctcca acgagtcctat catcgtcgac 1500
cgcagcaaca cgagtgtgc gccgagaaca actgctggga tcttcagtga taacgaggcg 1560
ggacgtctgc gctcttctga cgtgttgcga aatggaaaag aacaggttga aacittggag 1620
ctcactatcg tggtagataa tagtgtactg gaagtatatg ccaatggacg ctttgtctta 1680
ggcacttggg ctcggttaagt ctctcttggt tatggaagat tggtaaaaa ctaaccgcat 1740
gaaggcttgg gtacgccaac tcgactaaaa ttaacttctt ccataacggc gtggggagaag 1800
cgacattcga agatgtgacg gtctttgaag gactgtatga tgccctggcca caaaggaag 1859

<210> 8

<211> 22

<212> DNA

<213> Artificial Sequence

<400> 8

caactgctgg catcctcagt ga

<210> 9

<211> 29

<212> DNA

<213> Artificial Sequence

EP 1 022 332 A1

<400> 9

5 gcggatccat gaagctatca aatgcaatc

<210> 10

10 <211> 26

<212> DNA

<213> Artificial Sequence

15

<400> 10

20 gcggatcctt accgagccca agtgcc

<210> 11

25 <211> 27

<212> DNA

<213> Artificial Sequence

30

<400> 11

35 gcggatccctc acttcctttg tggccag

<210> 12

40 <211> 30

<212> DNA

<213> Artificial Sequence

45

<400> 12

50 gtggcgctac caagaccgag cccaagtgcc

55 <210> 13

EP 1 022 332 A1

<211> 1894

<212> DNA

<213> Scopulariopsis brevicaulis IF04843

<220>

<221> CDS

<222> (1)... (1721)

<221> intron

<222> (1722)... (1776)

<221> CDS

<222> (1777)... (1894)

<400> 13

caacctacgt ctcgtcaat cgacaattcc acgtatcctt ctatcgacta caactccgcc	60
cctccaaacc tctcgactct tgccaacaac agcctcttcg agacatggag gccgagggca	120
cacgtccttc cgccccagaa ccagatcggc gatccgigtg tgcactacac cgaccccgag	180
acaggaatct tccacgtcgg ctggctgtac aacggcaatg gcgcttcgg cgccacgacc	240
gaggatctcg tcacctatac ggatctcaac cccgacggag cgcagatgat ccttcgggt	300
ggtgtgaatg accccattgc tgcctttgac ggcgcggta ttcccagtgg catlgatggg	360
aaaccaccca tgatgtatac ctccggtgtc tacaigccca tctcctggag catcgcttac	420
accaggggaa gcgagaccca ctctctcgca gtgtcgtccg acggcggtaa gaacttcacc	480
aagctgggtc agggccccgt cattccttcg cctcccttcg gcgccaacgt gaccagctgg	540
cgtagccctt tctgttcca aaacccccag ttcgactctc tctctgaaag cgagaacggc	600
acgtgggtaca ccgttatctc tggteggcat cacggtgacg gccccccgc gtctctctac	660
cgtcagcacg accccgactt ccagtactgg gactaccttg gaccgtggig gaacgaggaa	720
gggaactcga cctggggcag cggtagctgg gctggccggt ggggctacaa cttegaggtc	780
atcaacattg tgggtcttga cgaatgatggc tacaaccccg acggtgaaat ctttgccacg	840
gtaggtaccg aatggtcgtt tgaccccatc aaaccgcagg cctcggacaa caggagatg	900
ctctgggccc cgggcaacat gactctcgag gacggcgata tcaagttcac gccaaagcatg	960
gcgggctacc tcgactgggg tctatcggcg tatgccgccc ctggcaagga gctgcccgct	1020

EP 1 022 332 A1

5
10
15
20
25
30

tcttcaaagc cttcgagaa gagcgggtgc cggaccggt tcgtgtcgta cctgtggctc 1080
accggtgact acttcgaggg ccacgacttc cccacccgc agcagaattg gaccggctcg 1140
cttttgcttc cgcgtgagct gagcgtcggg acgattccca acgttgtcga caacgagctt 1200
gctcgcgaga cgggctcttg gagggttggc accaacgaca ctggcgtgct tgagctggtc 1260
actctgaagc aggagattgc tcgcgagacg ctggctgaaa tgaccagcgg caactccttc 1320
accgaggcga gcaggaatgt cagctcgccc ggaictaccg ccttccagca gtccctggat 1380
tccaagtctt tcgtcctgac cgcctcgctc tcttccctt cgtcggctcg cgactccgac 1440
ctcaaggctg gtltcgagat cctgtcgtcc gagtttgagt cgaccacggt ctactaccag 1500
ttttccaacg agtccatcat cattgaccgg agcaactcga gtgctgccgc cttgactacc 1560
gatggaatcg acacccgcaa cgagtttggc aagatgcgcc tgtttgatgt tgtcgagggt 1620
gaccaggagc gtatcgagac gctcgatctc actattgtgg ttgataactc gatcgttgag 1680
gttcatgcc aaggcgatc cgctctgagc acttgggttc ggtaagtgga acgccgacca 1740
cgcttcttaa ttcttccaat actaacatta tcacagttcg tggtagagt cgtccaagga 1800
catcaagttc ttccacgatg gcgacagcac gggtcagttc tcgaacatca ccgtctacga 1860
gggactgttt gacgcctggc cggagcgggc cagg 1894

35
40

<210> 14

<211> 26

<212> DNA

<213> Artificial Sequence

45

<400> 14

gcggatccat gaaactctca actglt

50

<210> 15

<211> 26

<212> DNA

<213> Artificial Sequence

EP 1 022 332 A1

<400> 15**gcggatcctt accgaaccca agtgct****<210> 16****<211> 25****<212> DNA****<213> Artificial Sequence****<400> 16****gcggatcctt acctggcccg ctccg****<210> 17****<211> 33****<212> DNA****<213> Artificial Sequence****<400> 17****cgactcgtag cacgaacgaa cccaagtgct cag****Claims**

1. A polypeptide comprising the amino acid sequence of SEQ ID No. 1 or a homologue thereof.
2. A DNA encoding a polypeptide according to Claim 1.
3. A DNA according to Claim 2 comprising the nucleotide sequence of SEQ ID No. 2.
4. A polypeptide comprising the amino acid sequence of SEQ ID No. 3 or a homologue thereof.
5. A DNA encoding a polypeptide according to Claim 4.
6. A DNA according to Claim 5 comprising the nucleotide sequence of SEQ ID No. 4.
7. A vector comprising a DNA according to Claim 2, 3, 5 or 6.

EP 1 022 332 A1

8. A host cell transformed by a vector according to Claim 7.

9. A process for producing a β -fructofuranosidase comprising the steps of:

5 cultivating a host cell according to Claim 8, and
 collecting the β -fructofuranosidase from the host and/or the culture thereof.

10. A process for producing fructooligosaccharides comprising the step of bringing sucrose into contact with a host cell
10 according to Claim 8 or a β -fructofuranosidase obtained by the process according to Claim 9.

15

20

25

30

35

40

45

50

55

EP 1 022 332 A1




FIG. 1A

EP 1 022 332 A1




FIG. 1B

EP 1 022 332 A1




FIG. 1C

EP 1 022 332 A1

1.8 kbp BamHI DNA FRAGMENT B

pY2831

1.8 kbp BamHI DNA FRAGMENT A




FIG. 1D

EP 1 022 332 A1




FIG. 2 A

EP 1 022 332 A1




FIG. 2B

EP 1 022 332 A1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/04087

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl⁶ C12N9/10, C12N15/54, C12N1/19, C12P19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl⁶ C12N9/10, C12N15/54, C12N1/19, C12P19/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
DDBJ/EMBL/GenBank

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	L.M. Boddy et al., "Purification and characterization of an <i>Aspergillus niger</i> invertase and its DNA sequence", Curr. Genet., Vol. 24, P.60-66 (1993)	1-10
P, X	WO, 97/34004, A (Meiji Seika Kaisha, Ltd.), 18 September, 1997 (18. 09. 97) (Family: none)	1-10

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search
27 November, 1998 (27. 11. 98)Date of mailing of the international search report
8 December, 1998 (08. 12. 98)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.